

RESEARCH

Open Access



# Sarcopenia and its association with objectively measured life-space mobility and moderate-to-vigorous physical activity in the oldest-old amid the COVID-19 pandemic when a physical distancing policy is in force

Rick Yiu Cho Kwan<sup>1</sup>, Justina Yat Wa Liu<sup>2,3\*</sup>, Yue-Heng Yin<sup>2</sup>, Paul Hong Lee<sup>4</sup>, Siu Ying Ng<sup>5</sup>, Daphne Sze Ki Cheung<sup>2,3</sup>, Patrick Pui Kin Kor<sup>2</sup>, Simon Ching Lam<sup>1</sup>, Shirley Ka Lai Lo<sup>2</sup>, Lin Yang<sup>6</sup>, Siu Kay Chan<sup>5</sup> and Vico Chung Lim Chiang<sup>6</sup>

## Abstract

**Introduction:** The oldest-old are highly vulnerable to sarcopenia. Physical distancing remains a common and effective infection-control policy to minimize the risk of COVID-19 transmission during the pandemic. Sarcopenia is known to be associated with impaired immunity. Moderate-to-vigorous physical activity (MVPA) and life-space mobility (LSM) are potential strategies for minimizing the risk of sarcopenia. However, a physical distancing policy might jeopardize the practice of MVPA and LSM. The purposes of this study were to identify the prevalence of sarcopenia and examine the association between MVPA and LSM with sarcopenia in the community-dwelling oldest-old during the COVID-19 pandemic.

**Methods:** This study employed a cross-sectional and observational design. The study was conducted in 10 community centres for older people in Hong Kong during the period of the COVID-19 pandemic (September to December 2020). Eligible participants were the oldest-old people aged  $\geq 85$  years, who were community-dwelling and had no overt symptoms of cognitive impairment or depression. Key variables included sarcopenia as measured by SARC-F, LSM as measured by a GPS built into smartphones, and MVPA as measured by a wrist-worn ActiGraph GT3X+. Variables were described by mean and frequency. A multiple linear regression was employed to test the hypotheses. The dependent variable was sarcopenia and the independent variables included LSM and MVPA.

**Results:** This study recruited 151 eligible participants. Their mean age was 89.8 years and the majority of them were female ( $n = 93/151$ , 61.6%). The prevalence of sarcopenia was 24.5% ( $n = 37/151$ ) with a margin of error of 6.86%. MVPA was negatively associated with sarcopenia in older people ( $\beta = -0.002$ ,  $SE = 0.001$ ,  $p = 0.029$ ). However, LSM was not associated with sarcopenia.

\*Correspondence: justina.liu@polyu.edu.hk

<sup>3</sup> Research Institute for Smart Ageing, The Hong Kong Polytechnic University, Hung Hom, Hong Kong

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Conclusion:** The prevalence of sarcopenia in the community-dwelling oldest-old population is high. MVPA is negatively associated with sarcopenia. LSM is unrelated to sarcopenia. Sarcopenia should be recognized and the oldest-old with sarcopenia should be accorded priority treatment during the COVID-19 pandemic.

**Keywords:** Sarcopenia, Moderate-to-vigorous physical activity, Life-space mobility, COVID-19, Physical distancing

## Introduction

Sarcopenia is defined by the European Working Group on Sarcopenia in Older People (EWGSOP) as a reduction in appendicular skeletal muscle mass, muscle strength, and physical performance [1]. The prevalence of sarcopenia in older people ranges from 0.36 to 8.14%, varying according to the diagnostic criteria [2]. Sarcopenia is a disease linked to other diseases; thus, its prevalence is much higher in older people with chronic illnesses; for example, 42% of people with stroke, 31.4% of people with cardiovascular diseases, and 26.4% of people with dementia have sarcopenia [3, 4]. Recent evidence shows that older age is associated with a higher risk of developing sarcopenia and that the oldest-old with sarcopenia have higher odds of experiencing difficulties with physical function (e.g., stooping, kneeling, crouching, walking) [5]. Sarcopenia is also associated with a higher risk of many negative health outcomes, including death from all causes, in-hospital mortality, and falls [6].

Many countries have adopted physical distancing and stay-at-home policies as key strategies to slow the spread of the virus since the World Health Organization (WHO) declared coronavirus disease 2019 (COVID-19) to be a global pandemic in March 2020 [7, 8]. As in other regions during this pandemic, the Hong Kong Government adopted a physical distancing policy, advising the public to reduce their level of social contact. Specifically, people were instructed to maintain a distance of at least 1 m from others; minimize gatherings, group-based activities (e.g., fitness classes), and trips outside the home (particularly meal gatherings); avoid crowded places and physical contact (e.g., handshakes); and wear face masks [9]. Hong Kong people have shown good compliance with these infection-control measures [10, 11]. However, the prolonged implementation of the physical distancing and stay-at-home policies could increase the risk of people of all ages developing a variety of health problems [12]. Research has shown that older people with chronic conditions greatly decreased their physical activity levels and were more likely to be hospitalized during the pandemic [13, 14]. Older people, particularly the oldest-old (i.e., those aged over 85 years), may be more susceptible to the negative consequences of rigorous distancing [15].

It has been recommended that older people engage in at least 150 min of moderate-to-vigorous physical activity (MVPA), at 10-min bouts each time for beneficial health

outcomes [16]. MVPA is associated with many health benefits (e.g., reduced mortality) [17]. A significant reduction in physical activity levels worldwide has been identified as a negative impact of physical distancing [18]. A study showed that 2 weeks of inactivity could result in decreased muscle mass strength in approximately 8% of people, leading to ineffective muscle function [19]. An abrupt reduction in physical activity because of physical distancing is a particular concern for older people, [20] as all of these physiological changes may predispose older people to frailty and sarcopenia [21].

Life-space mobility (LSM) refers to movement extending from within one's household to beyond one's district or town [22]. Previous studies have demonstrated an association between declining life-space and an increased risk of frailty, [23] reduced physical activity, [24] and poor mental health [25]. A few studies have reported a significant reduction in LSM among older people during the COVID-19 pandemic [26, 27]. Studies conducted in Brazil and Finland demonstrated that restrictions in life-space during the COVID-19 pandemic had a negative impact on the quality of life of older people [15, 28].

Physical activity and engagement in activities outside the home are recommended treatments for sarcopenia in older people [29]. However, the LSM measured in studies of sarcopenia in older people has commonly been evaluated using questionnaires such as Life-Space Assessment [22]. Accuracy affected by the recall bias of participants is the major limitation of using a questionnaire to measure LSM [30]. Furthermore, there is a dearth of studies examining the associations among sarcopenia, physical activity, and LSM during the COVID-19 pandemic when strict infection-control policies (e.g., physical distancing, a stay-at-home policy) were in force; not to mention studies focusing on the most vulnerable older people (i.e., the oldest-old). This study will inform policymakers on how to sustain health in the oldest-old during the COVID-19 pandemic.

## Objectives

This study aims to:

1. Estimate the prevalence of sarcopenia in the oldest-old during the COVID-19 pandemic, and
2. Examine the association between sarcopenia with objectively measured MVPA and LSM.

## Methods

### Study design

This study employed a cross-sectional and observational design. To ensure clarity of reporting, we followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines [31].

### Setting

The study was conducted in ten community centres for older people in Hong Kong. The centres provide a range of activities for community-dwelling people aged  $\geq 60$  years to enable them to remain in the community and to lead healthy and dignified lives [32]. We adopted a convenience sampling method to invite community centres to join the study. The staff members in the community centres were briefed on the sample selection criteria. They then invited those of their members who met the criteria to be screened by our research team. The study, including recruitment and data collection, was conducted from September to December 2020. During this period, Hong Kong was experiencing the COVID-19 pandemic, with a mean daily number of 26 new cases (range = 0–115, median = 8) in a city with a population of approximately 7.5 million [33].

### Participants

Inclusion criteria were 1) the oldest-old people aged  $\geq 85$  years, [34] and 2) community-dwelling, defined as not having lived in a long-term care setting (e.g., nursing home) in the past 6 months before the data collection process. Participants were excluded if they were in a condition that could hinder them from providing accurate information, including 1) cognitive impairment, defined by a Mini-Cognitive Test score of  $\geq 3$ , [35] and depressive symptoms, defined by a Geriatric Depression Scale score of  $\geq 8$  [36, 37].

We asked staff members of the community centres to invite all potentially eligible participants to join the study through face-to-face and telephone invitations. Potentially eligible participants were then invited to attend the eligibility screening sessions, which were conducted in the community centres.

### Variables

Demographic variables included age, gender, marital status, educational attainment, and living conditions. Self-reporting was used to collect the demographic information of the participants, such as their living conditions, including type of residence and with whom they were living; and level of education, including the highest level of education that they had completed. Clinical variables included chronic illnesses, body mass index (BMI), and cognitive function. Predictor variables included life

space and MVPA. Outcome variables included sarcopenia. Data collection was conducted at the community centres by a trained research assistant through face-to-face interviews using validated instruments.

### Measurement


Chronic illnesses were measured using a questionnaire with six dichotomous questions (0 = no, 1 = yes) on common chronic illnesses, including Diabetes Mellitus, cardiovascular diseases, osteoarthritis, cancer, and hypertension, which were found to be associated with sarcopenia [4, 38].

Body mass index (BMI) was measured using a calibrated scale balance and a tape ruler employing the standard eq. ( $BMI = \text{body weight in kg}^2 / \text{body height in metres}$ ). BMI was categorized into four levels adjusted for Asian adults, as recommended by the WHO Western Pacific Region (underweight:  $BMI < 18.5$ , normal:  $BMI = 18.5\text{--}22.9$ , overweight:  $BMI = 23\text{--}25$ , and obese:  $BMI > 25$ ) [39].

Cognitive function was measured using the Abbreviated Mental Test (AMT) Hong Kong version [40]. AMT consists of 10 items evaluating orientation to time and place, attention, calculation, and memory. One point is assigned for a correct answer to each question, with a total score ranging from 0 to 10. A higher score indicates better cognitive function. It has been validated to have good internal consistency ( $\alpha = 0.814$ ), test-retest reliability ( $r = 0.993$ ), and concurrent validity with the Chinese Mini-Mental Stage Examination ( $r = 0.86$ ) [40, 41].

Sarcopenia was measured using the SARC-F questionnaire [42, 43]. SARC-F comprises five items assessing strength, assistance in walking, rising from a chair, climbing stairs, and falls. Each item was rated on a 3-point frequency scale (i.e., 0 = none to 2 = A lot) with a total score ranging from 0 to 10. A higher score indicates a higher risk of sarcopenia. It has been validated against the gold standard criteria of EWGSOP to have good validity in identifying sarcopenia (accuracy = 0.90–0.87) [43].

In this study, LSM is conceptualized as the area through which a person has travelled over a specific period, expanding from within one's home to beyond one's town or geographic region [44]. Therefore, LSM was measured using an objective measurement employing an Android-based smartphone (i.e., Redmi 9, Xiaomi) installed with a global positioning system (GPS) logging app. The location of the participants was collected using the Application Programming Interface (APK) provided by Google Maps. Data collection was automatically triggered every 15 min using Android Job Scheduler. As shown in Fig. 1, the participants were instructed to carry the smartphones with them continuously using the pouches provided. GPS is becoming more ubiquitous every day, as



most smartphones have a built-in GPS function. GPS uses a person’s geolocation to calculate various measures related to LSM, such as frequency and distance travelled from home. Some studies using GPS to determine LSM have reported promising results [45, 46]. The acceptability and feasibility of having older people use GPS has also been found to be satisfactory [47, 48]. Participants were asked to carry the smartphone with GPS at all times for 7 days.

GPS coordinates were categorized into five zones following the conceptualization of the Life Space Assessment (LSA) instrument validated in a Chinese older population [49]. Categorization of the coordinates into the five zones depended on the distance of the coordinates from the residential addresses of individual participants [50]. Given that GPS could not accurately differentiate between a bedroom and an apartment, we collapsed the zones of the bedroom and within the apartment building. We adopted a 4-concentric-circle categorization, with the centres at the participants’ residential address (i.e., zone 1=bedroom/within the apartment building, zone 2=neighbourhood other than own yard or apartment building, zone 3=outside the neighbourhood but within the town, and zone 4=outside the town). Based on a previous study, zone 1 was defined as the area of the concentric circular zone with a radius of 0–150m, while zone 4 was defined as the area beyond the circular zone with a radius of 1000m [50]. The cut-off points for zones 2 and 3 were estimated as being half of the difference between the radii of zones 1 and 4 (i.e., [150–1000m]/2). Thus, zones 2 and 3 were the circular

areas with radii of 51–580m and 581–1000m, respectively. The percentage of time spent in each zone over the total amount of time spent in collecting data was calculated. An LSM score was computed by summing the weighted percentage of time spent in each zone. The weighting was assigned according to the distance of the zone from the participant’s home (i.e., zone 1 = 1.5, zone 2 = 3, zone 3 = 4, and zone 4 = 5). The weighting assigned to each zone refers to the weighting used in the Life Space Questionnaire [49]. A higher LSM score indicates a higher level of LSM. Details regarding the smartphone wearing compliance and GPS logging methods could be found in the [Supplementary Materials](#).

MVPA was measured using a wrist-worn ActiGraph GT3X+, which was mounted on the participants’ non-dominant wrist for 24h for 7 days. ActiGraph GT3X+ was validated against indirect calorimetry to accurately identify MVPA (sens = 0.836, spec = 0.894, AUC = 0.932). Participants were instructed to wear the device continuously during the assessment period [51]. An MVPA minute was defined as a minute in which the ActiGraph recorded physical movement (i.e., vector magnitude) of above 4117.7 cpm [51]. Only at least 10min of continuous MVPA were counted as valid minutes because only such sessions are considered beneficial by the WHO [16]. MVPA minutes were measured over seven consecutive days with an epoch length of 1 minute [52]. Participants were instructed to wear the ActiGraph continuously throughout the 7-day study period (i.e., 24h/day for 7 days). We used 60 min of a continuous count of zero to determine “non-wear time” [53]. Only MVPA minutes

measured on valid days (i.e., ActiGraph wear time > 10h/day) for a valid period (i.e., valid days  $\geq 4$  days) were entered into the data analysis [54, 55].

### Study size

To estimate the sample size needed to determine the prevalence of sarcopenia in the oldest-old (i.e., objective #1), Cochran's formula was employed [56]. We assumed that the prevalence of sarcopenia was 7.3%, as observed in a local study of community-dwelling older people in Hong Kong [57]. Cochran's formula showed that 163 participants were needed, with a confidence interval of 95% and a margin of error of 4% in the population of the oldest-old (i.e., people aged  $\geq 85$  years) of approximately 170,000 people, as estimated in the 2016 Hong Kong census [57].

To estimate the sample size needed to determine the association between sarcopenia and MVPA minutes and LSM, the following linear multiple regression was employed: Fixed model,  $R^2$  increase package in G\*Power. The effect size of accelerometer-determined MVPA minutes on sarcopenia was small-medium (i.e., Cohen's  $d=0.47$ ) [58, 59]. We did not consider the effect size of LSM on sarcopenia because this information was not available in the literature. G\*Power showed that 158 participants would be needed, with the parameters estimated to be  $f^2=0.1$ ,  $\alpha=0.05$ , power=0.95, number of tested predictors = 2, and the total number of predictors = 14.

To ensure an adequate number of samples to fulfil the two research objectives, we adopted a sample size of 163.

### Statistical methods

Demographic variables, clinical variables, and outcomes were described using mean and frequency according to their levels of measurement. To test the association between sarcopenia with LSM and physical activity (i.e., objective #2), multiple linear regression was employed. The dependent variable was sarcopenia. The independent variables were LSM and MVPA. Potential confounding variables included age, gender, education, chronic diseases, cognitive function, living condition, and BMI because they are known to be associated with sarcopenia, as well as LSM and physical activity [6, 28].

Missing data on the demographic and clinical variables were replaced by mean values. Missing MVPA data were defined as those detected as non-wearing, and missing GPS data were defined as data collected within an invalid period or collected with poor accuracy. Both GPS and MVPA data were replaced by mean values, and only available data on valid days were entered into the data analysis [54, 55]. A sensitivity analysis comparing results between two missing data management strategies (i.e., replacement by mean) was performed.

## Results

### Participants

As shown in Fig. 1, we recruited participants in 10 community centres for older people. We assessed the eligibility of 189 members of these centres. In the pre-screening phase, 10,976 members were identified as not eligible to take part in the study, through a reading of their records in the centres by centre staff. Common reasons for exclusion included age, a documented condition of dementia, and an inability to speak Cantonese. We identified 163 eligible participants, but 11 did not consent to data collection and 1 did not complete the data collection process. In the end, we included data from 151 participants for analysis.

There were no missing data on the demographic and clinical variables, except on BMI, which was replaced by the mean value. There were 32 invalid cases (21.2%) of Actigraph data and 26 invalid cases (17.2%) of GPS data; these were replaced by the mean values of other participants. There were no significant differences in missing data between the groups with and without sarcopenia.

### Descriptive data

As shown in Table 1, the participants' mean age was 89.8 years. The majority were female ( $n=96$ , 63.6%), widowed/divorced/separated ( $n=93$ , 61.6%), uneducated ( $n=79$ , 52.3%), living with family/domestic helper/friends ( $n=88$ , 57.9%), had hypertension ( $n=108$ , 71.5%), and normal BMI ( $n=61$ , 40.4%). The mean AMT score was 8.5 (SD=1.3). The mean SARC-F score was 2.1 (SD=2.1). The mean number of MVPA minutes per week was 181.3 (SD=202.5). The mean LSM score was 1035.3 (SD=141.7). The mean percentage of time spent in zone 1 was 84.0% (SD=13.0), in zone 2 was 12.6% (SD=12.0), in zone 3 was 1.2% (SD=2.4), and in zone 4 was 2.3% (SD=4.6) (Fig. 2).

### Main results

#### Objective #1

The prevalence of sarcopenia was 24.5% ( $n=37/151$ ) with a margin of error of 6.86%.

#### Objective #2

As shown in Table 2, MVPA was negatively associated with sarcopenia in older people ( $\beta = -0.002$ , SE = 0.001) and the association was statistically significant ( $p=0.047$ ). However, the LSM score was not associated with sarcopenia. Age and cognitive function were also observed to be associated with sarcopenia. These factors explained 27% of the variance for sarcopenia ( $R^2=0.270$ ). A sensitivity analysis did not reveal any differences in

**Table 1** Participants' demographic and clinical profile, predictors, and outcomes ( $N = 151$ )

<b>Demographic and clinical profile</b>	<b>mean (SD) / n (%)</b>
Age, mean year (SD)	89.8 (3.6)
Gender, n (%)	
Male	55 (36.2)
Female	96 (63.6)
Marital status, n (%)	
Unmarried	2 (1.3)
Married	56 (36.8)
Widowed/divorced/separated	93 (61.6)
Educational attainment, n (%)	
Tertiary	2 (1.3)
Secondary	13 (8.6)
Primary	63 (41.7)
Nil	73 (48.3)
Living condition, n (%)	
Living alone	63 (41.7)
Living with family/domestic helper/friends	88 (57.9)
Chronic diseases, n (%)	
Diabetes Mellitus	21 (13.9)
Heart diseases	26 (17.2)
Osteoarthritis	8 (5.3)
Cancer	5 (3.3)
Hypertension	108 (71.5)
BMI, mean (SD)	23.6 (3.7)
BMI, n (%)	
Underweight (BMI < 18.5)	8 (5.3)
Normal (BMI = 18.5–22.9)	61 (40.4)
Overweight (BMI = 23–25)	36 (23.8)
Obese (BMI > 25)	46 (30.5)
Cognitive function	
AMT, mean (SD)	8.5 (1.3)
<i>Predictors</i>	
MVPA	
Minutes per week, mean (SD)	181.3 (202.5)
Life-space mobility score, mean (SD)	1035.3 (141.7)
Time spent in each life-space zone, mean % (SD)	
Zone 1	84.0 (13.0)
Zone 2	12.6 (12.0)
Zone 3	1.2 (2.4)
Zone 4	2.3 (4.6)
<i>Outcomes</i>	
Sarcopenia (SARC-F), mean (SD)	2.1 (2.1)
Sarcopenia (SARC-F $\geq 4$ ), n (%)	
Yes	37 (24.5)
No	114 (75.5)

*BMI* body mass index, *AMT* Abbreviated Mental Test, *MVPA* moderate-to-vigorous physical activity, *Zone 1* bedroom/within the apartment building, *Zone 2* neighbourhood other than own yard or apartment building, *Zone 3* outside neighbourhood but within the town, *Zone 4* outside the town, *SARC-F* SARC-F questionnaire

the interpretation of the results between the two missing data management strategies (i.e., list-wise deletion and replacement by mean).

## Discussion

This appears to be the first study to examine the association between sarcopenia with objectively measured MVPA and LSM in community-dwelling oldest-old people. There are three key findings. First, the prevalence of sarcopenia in the oldest-old during the COVID-19 pandemic was high. Second, MVPA was negatively associated with sarcopenia in the oldest-old. Third, the MVPA of the majority of the oldest-old was adequate (i.e., MVPA min > 150 min/week) even though the participants did not go far from their residence. These findings have several implications.

A study that surveyed 670 older people in the Chinese community with a mean age of 76.2 years, and employing the same method (i.e., SARC-F score  $\geq 4$ ), showed that the prevalence of sarcopenia was 6.1% [60]. A systematic review of seven studies ( $n = 12,800$ ) that included older people with a mean age of 75.1 years and that employed various diagnostic criteria (e.g., EWGSOP, International Working group on Sarcopenia, Asian Working Group for Sarcopenia, and Foundation for the National Institutes of Health Sarcopenia Project) showed that the prevalence of sarcopenia in older people was 3–17.5% [61]. A study examining the prevalence of sarcopenia specifically in the oldest-old community-dwelling Chinese showed that the prevalence increased dramatically from 15.1% in those aged 80–84 years to 63.6% in those aged 95+ years [5]. In comparison, the percentage of older people with sarcopenia observed in this study was lower (i.e., 24.5%). A possible reason for why more older people in our study were observed to have sarcopenia is that the effects of COVID-19 (e.g., confinement on physical activity, dietary habits, sleep, and stress) might have given rise to an increased likelihood of muscle loss [21]. The high prevalence of sarcopenia could also be due to the age of the sample in this study, which included only the oldest-old. The prevalence of sarcopenia is known to increase exponentially with age [5]. This study suggests that COVID-19 prevention measures (e.g., lockdown) might have led to more nutrition problems, reduced physical activity, and increased social isolation, all of which would lead to sarcopenia. The oldest-old are a particularly vulnerable segment of the population. Sarcopenia could subsequently induce impaired immune function and heightened metabolic stress, [62] leading to susceptibility to COVID-19 infection. Therefore, this study recommends that sarcopenia be recognized and treated as a priority during the COVID-19 pandemic.



This study adopted the approach of including all forms of MVPA that last continuously for 10 min in a free-living setting. Given that it could be difficult for older people in free-living settings to recognize MVPA, it is recommended that they wear trackers to help to ensure that they attain an intensity consistent with the definition of MVPA when they exercise. This is because a systematic review showed that commercially available wearable trackers are valid for measuring MVPA [63]. Therefore, this study recommends that the oldest-old practise MVPA in any form for 10 continuous minutes gauged by wearable trackers to combat sarcopenia during the COVID-19 period. For example, brisk walking in a free-living setting could be intense enough to achieve MVPA and could feasibly be practised by older people in free-living settings [64].

A recent study conducted in a Chinese population with a mean age of 70 years before the COVID-19 pandemic showed that older people spent an average of 175.0 min in MVPA per week [65]. This is very similar to what we observed in this study (i.e., 181.3 min/week). This shows that the amount of MVPA practised by the oldest-old is similar to the average for older people and has been unaffected by the COVID-19 pandemic. The average amount of MVPA in the oldest-old is above the beneficial amount of 150 min/week as recommended by the WHO. The GPS data indicated that the vast majority of participants did not go far from their apartments. MVPA is unlikely to be achieved at home given the small living spaces in Hong Kong. This study showed that it was feasible to remain physically active without going far from home. Also, despite the adequate amount of MVPA on average, the

prevalence of sarcopenia was still somewhat high. Future studies should examine what other factors contributed to the higher level of sarcopenia in the oldest-old during the COVID-19 pandemic. This study, therefore, recommends that the oldest-old remain physically active (i.e., MVPA > 150 min/week) during the COVID-19 pandemic, pay attention to other risk factors of sarcopenia apart from inadequate physical activity (e.g., poor nutrition), and avoid unnecessary travel to minimize the risk of contracting COVID-19 [66].

This study has several limitations. First, the sample size was relatively small. Caution should be exercised on the claim of the prevalence of sarcopenia because the margin of error is not low (i.e., 6.86%). Second, although we had already conducted subject screening and recruitment at 10 community centres, those were convenience samples and no sampling frame was used. Caution should also be exercised on the issue of the generalizability of the findings to the population. Third, there is a notable amount of missing data relating to the MVPA and GPS. Yet a sensitivity analysis showed no differences in interpretation of the results between the two management methods (i.e., replaced by mean values and listwise deletion). Fourth, because of the short battery life of the smartphone, GPS data could not be mandatorily sampled every 15 min. Instead, the GPS data copied the previous coordinate every 15 min if no movement was detected. The “no movement” assumption could also be a case of poor reception of GPS signals. Fifth, the effect size of MVPA on sarcopenia was observed to be very strong, and significant. Sixth, the SARC-F has low sensitivity, so we might have dismissed positive cases, meaning that the actual

**Table 2** Regression model predicting sarcopenia by SARC-F score

<i>N</i> = 151, <i>R</i> <sup>2</sup> = 0.295	Beta (SE)	<i>p</i> -value	95%CI
<i>Confounding factors</i>			
Gender			
Female	1.573 (0.403)	< 0.001*	0.776, 2.371
Male	0		
Age	0.017 (0.049)	0.733	−0.080, 0.114
Education			
Tertiary	−1.820 (1.395)	0.149	−4.579, 0.940
Secondary	−0.232 (0.618)	0.708	−1.454, 0.990
Primary	−0.290 (0.359)	0.421	−1.002, 0.421
Nil	0		
Cognitive function (AMT)	−0.333 (0.133)	0.013*	−0.595, −0.070
Marital		0.141	−0.055, 0.380
Unmarried	0.432 (1.395)	0.757	−2.328, 3.192
Married	0.074 (0.409)	0.856	−0.735, 0.883
Widowed/divorced/separated	0		
Living condition			
Living alone	−0.391 (0.352)	0.270	−1.088, 0.307
Living with family/domestic helper/friends	0		
BMI	0.062 (0.045)	0.177	−0.028, 0.151
Chronic diseases			
Diabetes Mellitus	−0.019 (0.476)	0.968	−0.961, 0.923
Heart diseases	0.045 (0.425)	0.915	−0.795, 0.886
Osteoarthritis	0.783 (0.702)	0.267	−0.606, 2.172
Cancer	0.117, (0.894)	0.896	−1.652, 1.887
Hypertension	−0.118 (0.375)	0.753	−0.860, 0.623
<i>Predictors</i>			
MVPA minutes per week	−0.002 (0.001)	0.047*	−0.003, −2.572
Life Space Mobility	0.038 (0.052)	0.464	−0.064, 0.140
Time spent in each life-space zone			
Zone 1	44.499 (67.279)	0.509	−88.568, 177.566
Zone 2	6.456 (44.623)	0.877	−75.866, 88.778
Zone 3	−21.628 (44.954)	0.631	−110.539, 67.282
Zone 4	−40.724 (62.422)	0.515	−164.184, 82.735

prevalence of sarcopenia might be higher. But the SARC-F is still an effective tool for screening potential cases of sarcopenia because of its high specificity (relatively good overall diagnostic accuracy) [43]. Finally, there is the possibility of an undetected residual confounding bias, so the findings need to be treated dialectically.

## Conclusion

The prevalence of sarcopenia in the community-dwelling oldest-old population is high. MVPA is negatively associated with sarcopenia. LSM is unrelated to sarcopenia. During the COVID-19 pandemic, sarcopenia should be recognized and treated as a priority for the oldest-old.

## Abbreviations

COVID-19: Coronavirus Disease of 2019; MVPA: Moderate-to-Vigorous Physical Activity; LSM: Life-Space Mobility; SARC-F: Strength, Assistance with walking, Rising from a chair, Climbing stairs, and Falls; EWGSOP: European Working Group on Sarcopenia in Older People; WHO: World Health Organization; STROBE: Strengthening the Reporting of Observational Studies in Epidemiology; BMI: Body Mass Index; AMT: Abbreviated Mental Test; GPS: Global Positioning System; APK: Android Application Package; AGPS: Assisted Global Positioning System; GLONASS: Global Navigation Satellite System.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12877-022-02861-7>.

Additional file 1.



### Acknowledgements

We thank all our participants for their contributions to this study. We also thank Ms. Lydia Suen for formatting the content and responses.

### Authors' contributions

R.K., J.L., P.L., S.N., D.C., P.K., S.L., L.Y., S.C., and V.C. discussed and developed the idea for the study. J.L. and S.N., coordinated the collecting of data. P.L. designed the statistical plan and wrote up the statistical analysis plan. J.L., D.C., P.K., S.L., L.Y., S.C., and V.C. wrote the introduction. R.K. wrote the methods, results, and discussion sections. Y.Y.H revised the manuscript. All of the authors contributed equally to critically reviewing and commenting on the manuscript. The author(s) read and approved the final manuscript.

### Funding

The project was supported by Sik Sik Yuen (Ref: ZH3X), a non-governmental organization in Hong Kong that provides social services to older people in Hong Kong. Apart from introducing potential participants for the study, the funder did not engage in any other part of the research for the study nor in the writing of the manuscript.

### Availability of data and materials

The datasets generated and analysed during the current study are not publicly available due to the restrictions involved when obtaining ethical approval for our study, which commit us to sharing the data only with members of the research team, but allow data to be made available from the corresponding author upon reasonable request.

### Declarations

#### Ethics approval and consent to participate

The study was approved by the Institutional Review Board of The Hong Kong Polytechnic University (Ref no.: HSEARS20200722001). The study was explained to all of the participants, and all gave their signed informed consent to take part in this study. Written informed consent and signatures were obtained from all participants in the study and for illiterate participants from their legal guardian/LAR. The study was conducted in accordance with the principles of the Declaration of Helsinki.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare that they have no competing interests.

#### Author details

<sup>1</sup>School of Nursing, Tung Wah College, Sheung Wan, Hong Kong. <sup>2</sup>Centre for Gerontological Nursing, School of Nursing, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong. <sup>3</sup>Research Institute for Smart Ageing, The Hong Kong Polytechnic University, Hung Hom, Hong Kong. <sup>4</sup>Department of Health Sciences, University of Leicester, Leicester, UK. <sup>5</sup>Sik Sik Yuen, Kowloon, Hong Kong. <sup>6</sup>School of Nursing, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong.

Received: 27 July 2021 Accepted: 16 February 2022

Published online: 25 March 2022

### References

- Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*. 2018;48(1):16–31. <https://doi.org/10.1093/ageing/afy169>.
- Petermann-Rocha F, Chen M, Gray SR, Ho FK, Pell JP, Celis-Morales C. New versus old guidelines for sarcopenia classification: What is the impact on prevalence and health outcomes? *Age Ageing*. 2019;49(2):300–4. <https://doi.org/10.1093/ageing/afz126>.
- Su Y, Yuki M, Otsuki M. Prevalence of stroke-related sarcopenia: a systematic review and meta-analysis. *J Stroke Cerebrovasc Dis*. 2020;29(9):105092. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2020.105092>.
- Pacifico J, Geerlings MAJ, Reijnierse EM, Phassouliotis C, Lim WK, Maier AB. Prevalence of sarcopenia as a comorbid disease: a systematic review and meta-analysis. *Exp Gerontol*. 2020;131:110801. <https://doi.org/10.1016/j.exger.2019.110801>.
- Xu W, Chen T, Cai Y, Hu Y, Fan L, Wu C. Sarcopenia in community-dwelling oldest old is associated with disability and poor physical function. *J Nutr Health Aging*. 2020;24(3):339–45. <https://doi.org/10.1007/s12603-020-1325-4>.
- Marzetti E, Calvani R, Tosato M, Cesari M, Di Bari M, Cherubini A, et al. Sarcopenia: an overview. *Aging Clin Exp Res*. 2017;29(1):11–7. <https://doi.org/10.1007/s40520-016-0704-5>.
- World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020 [updated 11 March 2020. Available from: <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19-20200311>.
- Lewnard JA, Lo NC. Scientific and ethical basis for social-distancing interventions against COVID-19. *Lancet Infect Dis*. 2020;20(6):631–3. [https://doi.org/10.1016/s1473-3099\(20\)30190-0](https://doi.org/10.1016/s1473-3099(20)30190-0).
- The Government of the Hong Kong Special Administrative Region. Social Distancing 2021. Available from: [https://www.coronavirus.gov.hk/eng/social\\_distancing.html](https://www.coronavirus.gov.hk/eng/social_distancing.html). [updated 2 Jul 2021].
- Zhao SZ, Wong JYH, Wu Y, Choi EPH, Wang MP, Lam TH. Social distancing compliance under COVID-19 pandemic and mental health impacts: a population-based study. *Int J Environ Res Public Health*. 2020;17(18). <https://doi.org/10.3390/ijerph17186692>.
- Bowman L, Kwok KO, Redd R, Yi Y, Ward H, Wei W, et al. Comparing public perceptions and preventive behaviors during the early phase of the COVID-19 pandemic in Hong Kong and the United Kingdom: cross-sectional survey study. *J Med Internet Res*. 2021;23(3):e23231. <https://doi.org/10.2196/23231>.
- Ford MB. Social distancing during the COVID-19 pandemic as a predictor of daily psychological, social, and health-related outcomes. *J Gen Psychol*. 2021;148(3):249–71. <https://doi.org/10.1080/00221309.2020.1860890>.
- López-Bueno R, Torres-Castro R, Koyanagi A, Smith L, Soysal P, Calatayud J. Associations between recently diagnosed conditions and hospitalization due to COVID-19 in patients aged 50 years and older: A SHARE-based analysis. *J Gerontol A Biol Sci Med Sci*. 2021. <https://doi.org/10.1093/geron/0/glab199>.
- López-Sánchez GF, López-Bueno R, Gil-Salmerón A, Zauder R, Skalska M, Jastrzębska J, et al. Comparison of physical activity levels in Spanish adults with chronic conditions before and during COVID-19 quarantine. *Eur J Pub Health*. 2021;31(1):161–6. <https://doi.org/10.1093/eurpub/ckaa159>.
- Saraiva MD, Apolinario D, Avelino-Silva TJ, de Assis Moura Tavares C, Gattás-Vernaglia IF, Marques Fernandes C, et al. The impact of frailty on the relationship between life-space mobility and quality of life in older adults during the COVID-19 pandemic. *J Nutr Health Aging*. 2021;25(4):440–7. <https://doi.org/10.1007/s12603-020-1532-z>.
- World Health Organization. Global Recommendations on Physical Activity for Health 2010.
- Hupin D, Roche F, Gremeaux V, Chatard J-C, Oriol M, Gaspoz J-M, et al. Even a low-dose of moderate-to-vigorous physical activity reduces mortality by 22% in adults aged ≥60 years: a systematic review and meta-analysis. *J Br J Sports Med*. 2015;49(19):1262–7. <https://doi.org/10.1136/bjsports-2014-094306>.
- Tison GH, Avram R, Kuhar P, Abreau S, Marcus GM, Pletcher MJ, et al. Worldwide effect of COVID-19 on physical activity: a descriptive study. *Ann Intern Med*. 2020;173(9):767–70. <https://doi.org/10.7326/m20-2665>.
- Reidy PT, McKenzie AI, Mahmassani Z, Morrow VR, Yonemura NM, Hopkins PN, et al. Skeletal muscle ceramides and relationship with insulin sensitivity after 2 weeks of simulated sedentary behaviour and recovery in healthy older adults. *J Physiol*. 2018;596(21):5217–36. <https://doi.org/10.1113/jp276798>.
- Roschel H, Artioli GG, Gualano B. Risk of increased physical inactivity during COVID-19 outbreak in older people: a call for actions. *J Am Geriatr Soc*. 2020;68(6):1126–8. <https://doi.org/10.1111/jgs.16550>.
- Kirwan R, McCullough D, Butler T, Perez de Heredia F, Davies IG, Stewart C. Sarcopenia during COVID-19 lockdown restrictions: long-term health effects of short-term muscle loss. *GeroScience*. 2020;42(6):1547–78. <https://doi.org/10.1007/s11357-020-00272-3>.

22. Peel C, Sawyer Baker P, Roth DL, Brown CJ, Brodner EV, Allman RM. Assessing mobility in older adults: the UAB study of aging life-space assessment. *Phys Ther.* 2005;85(10):1008–119.
23. Kwan RYC, Cheung DSK, Lo SKL, Ho LYW, Katigbak C, Chao YY, et al. Frailty and its association with the Mediterranean diet, life-space, and social participation in community-dwelling older people. *Geriatric Nurs (New York, NY).* 2019;40(3):320–6. <https://doi.org/10.1016/j.gerinurse.2018.12.011>.
24. Tsai LT, Rantakokko M, Rantanen T, Viljanen A, Kauppinen M, Portegijs E. Objectively measured physical activity and changes in life-space mobility among older people. *J Gerontol A Biol Sci Med Sci.* 2016;71(11):1466–71. <https://doi.org/10.1093/gerona/glw042>.
25. Byles JE, Leigh L, Vo K, Forder P, Curry C. Life space and mental health: a study of older community-dwelling persons in Australia. *Aging Ment Health.* 2015;19(2):98–106. <https://doi.org/10.1080/13607863.2014.917607>.
26. Perracini MR, de Amorim JSC, Lima CA, da Silva A, Trombini-Souza F, Pereira DS, et al. Impact of COVID-19 pandemic on life-space mobility of older adults living in Brazil: REMOBILIZE study. *Front Public Health.* 2021;9:643640. <https://doi.org/10.3389/fpubh.2021.643640>.
27. Rantanen T, Eronen J, Kauppinen M, Kokko K, Sanaslahti S, Kajan N, et al. Life-space mobility and active aging as factors underlying quality of life among older people before and during COVID-19 lockdown in Finland—a longitudinal study. *J Gerontol A Biol Sci Med Sci.* 2021;76(3):e60–e7. <https://doi.org/10.1093/gerona/glaa274>.
28. Rantakokko M, Iwarsson S, Portegijs E, Viljanen A, Rantanen T. Associations between environmental characteristics and life-space mobility in community-dwelling older people. *J Aging Health.* 2014;27(4):606–21. <https://doi.org/10.1177/0898264314555328>.
29. Uemura K, Makizako H, Lee S, Doi T, Lee S, Tsutsumimoto K, et al. The impact of sarcopenia on incident homebound status among community-dwelling older adults: a prospective cohort study. *Maturitas.* 2018;113:26–31. <https://doi.org/10.1016/j.maturitas.2018.03.007>.
30. Taylor JK, Buchan IE, van der Veer SN. Assessing life-space mobility for a more holistic view on wellbeing in geriatric research and clinical practice. *Aging Clin Exp Res.* 2019;31(4):439–45. <https://doi.org/10.1007/s40520-018-0999-5>.
31. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The strengthening of reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Int J Surg.* 2014;12(12):1495–9. <https://doi.org/10.1016/j.jisu.2014.07.013>.
32. Social Welfare Department of the Government of the Hong Kong Special Administrative Region. District Elderly Community Centre Hong Kong; 2021. Available from: [https://www.swd.gov.hk/en/index/site\\_pubsvc/page\\_elderly/sub\\_csselderly/id\\_districtel/](https://www.swd.gov.hk/en/index/site_pubsvc/page_elderly/sub_csselderly/id_districtel/).
33. Centre for Health Protection. 2020. Available from: <https://chp-dashboard.geodata.gov.hk/covid-19/en.html>. [updated 9 Dec 2020].
34. Smith J, Borchelt M, Maier H, Jopp D. Health and well-being in the young old and oldest old. *J Soc Issues.* 2002;58(4):715–32. <https://doi.org/10.1111/1540-4560.00286>.
35. Borson S, Scanlan JM, Chen P, Ganguli M. The Mini-cog as a screen for dementia: validation in a population-based sample. *J Am Geriatr Soc.* 2003;51(10):1451–4. <https://doi.org/10.1046/j.1532-5415.2003.51465.x>.
36. H-cB L, Chiu HFK, Kowk WY, Leung CM, et al. Chinese elderly and the GDS short form: a preliminary study. *Clinical gerontologist: the journal of Aging Ment Health.* 1993;14(2):37–42.
37. Leshner EL, Berryhill JS. Validation of the geriatric depression scale-short form among inpatients. *J Clin Psychol.* 1994;50(2):256–60. [https://onlinelibrary.wiley.com/doi/10.1002/1097-4679\(199403\)50:2%3C256::AID-JCLP2270500218%3E3.0.CO;2-E](https://onlinelibrary.wiley.com/doi/10.1002/1097-4679(199403)50:2%3C256::AID-JCLP2270500218%3E3.0.CO;2-E).
38. Veronese N, Punzi L, Sieber C, Bauer J, Reginster J-Y, Maggi S, et al. Sarcopenic osteoarthritis: a new entity in geriatric medicine? *Eur Geriatr Med.* 2018;9(2):141–8. <https://doi.org/10.1007/s41999-018-0034-6>.
39. Anuurad E, Shiwaku K, Nogi A, Kitajima K, Enkhmaa B, Shimono K, et al. The new BMI criteria for Asians by the regional Office for the Western Pacific Region of WHO are suitable for screening of overweight to prevent metabolic syndrome in elder Japanese workers. *J Occup Health.* 2003;45(6):335–43. <https://doi.org/10.1539/joh.45.335>.
40. Lam SC, Wong YY, Woo J. Reliability and validity of the abbreviated mental test (Hong Kong version) in residential care homes. *J Am Geriatr Soc.* 2010;58(11):2255–7. <https://doi.org/10.1111/j.1532-5415.2010.03129.x>.
41. Chiu HFK. Reliability and validity of the Cantonese version of mini-mental state examination—a preliminary study. *Hong Kong J Psychiatry.* 1994;4(2):25.
42. Malmstrom TK, Morley JE. SARC-F: a simple questionnaire to rapidly diagnose sarcopenia. *J Am Med Dir Assoc.* 2013;14(8):531–2.
43. Woo J, Leung J, Morley JE. Validating the SARC-F: a suitable community screening tool for sarcopenia? *J Am Med Dir Assoc.* 2014;15(9):630–4. <https://doi.org/10.1016/j.jamda.2014.04.021>.
44. May D, Nayak US, Isaacs B. The life-space diary: a measure of mobility in old people at home. *Int Rehabil Med.* 1985;7(4):182–6. <https://doi.org/10.3109/03790798509165993>.
45. Hirsch JA, Winters M, Clarke P, McKay H. Generating GPS activity spaces that shed light upon the mobility habits of older adults: a descriptive analysis. *Int J Health Geogr.* 2014;13:51. <https://doi.org/10.1186/1476-072x-13-51>.
46. Takemoto M, Carlson JA, Moran K, Godbole S, Crist K, Kerr J. Relationship between objectively measured transportation behaviors and health characteristics in older adults. *Int J Environ Res Public Health.* 2015;12(11):13923–37. <https://doi.org/10.3390/ijerph121113923>.
47. Webber SC, Porter MM. Monitoring mobility in older adults using global positioning system (GPS) watches and accelerometers: a feasibility study. *J Aging Phys Act.* 2009;17(4):455–67. <https://doi.org/10.1123/japa.17.4.455>.
48. Shoval N, Wahl H-W, Auslander G, Isaacson M, Oswald F, Edry T, et al. Use of the global positioning system to measure the out-of-home mobility of older adults with differing cognitive functioning. *J Ageing Soc.* 2011;31(5):849.
49. Ji M, Zhou Y, Liao J, Feng F. Pilot study on the Chinese version of the life space assessment among community-dwelling elderly. *Arch Gerontol Geriatr.* 2015;61(2):301–6. <https://doi.org/10.1016/j.archger.2015.06.012>.
50. Fillekes MP, Röcke C, Katana M, Weibel R. Self-reported versus GPS-derived indicators of daily mobility in a sample of healthy older adults. *Soc Sci Med.* 2019;220:193–202. <https://doi.org/10.1016/j.socscimed.2018.11.010>.
51. Kwan RYC, Liu JYW, Lee D, Tse CYA, Lee PH. A validation study of the use of smartphones and wrist-worn ActiGraphs to measure physical activity at different levels of intensity and step rates in older people. *Gait Posture.* 2020;82:306–12. <https://doi.org/10.1016/j.gaitpost.2020.09.022>.
52. Hart TL, Swartz AM, Cashin SE, Strath SJ. How many days of monitoring predict physical activity and sedentary behaviour in older adults? *Int J Behav Nutr Phys Act.* 2011;8(1):62. <https://doi.org/10.1186/1479-5868-8-62>.
53. Gorman E, Hanson HM, Yang PH, Khan KM, Liu-Ambrose T, Ashe MC. Accelerometry analysis of physical activity and sedentary behavior in older adults: a systematic review and data analysis. *Eur Rev Aging Phys Act.* 2014;11(1):35–49. <https://doi.org/10.1007/s11556-013-0132-x>.
54. Lee JA, Williams SM, Brown DD, Laurson KR. Concurrent validation of the Actigraph gt3x+, polar active accelerometer, Omron HJ-720 and Yamax Digiwalker SW-701 pedometer step counts in lab-based and free-living settings. *J Sports Sci.* 2015;33(10):991–1000. <https://doi.org/10.1080/02640414.2014.981848>.
55. Sasaki JE, John D, Freedson PS. Validation and comparison of ActiGraph activity monitors. *J Sci Med Sport.* 2011;14(5):411–6. <https://doi.org/10.1016/j.jsams.2011.04.003>.
56. Cochran WG. Sampling techniques. 3rd ed. New York: John Wiley; 1977.
57. Census and statistics department Hong Kong special Administrative region. Thematic report: older persons. In: Region CaSDHKSA, editor. Hong Kong; 2018.
58. Chinn S. A simple method for converting an odds ratio to effect size for use in meta-analysis. *Stat Med.* 2000;19(22):3127–31. [https://doi.org/10.1002/1097-0258\(20001130\)19:22<3127::aid-sim784>3.0.co;2-m](https://doi.org/10.1002/1097-0258(20001130)19:22<3127::aid-sim784>3.0.co;2-m).
59. Scott D, Johansson J, Gandham A, Ebeling PR, Nordstrom P, Nordstrom A. Associations of accelerometer-determined physical activity and sedentary behavior with sarcopenia and incident falls over 12 months in community-dwelling Swedish older adults. *J Sport Health Sci.* 2020. <https://doi.org/10.1016/j.jshs.2020.01.006>.
60. Wu T-Y, Liaw C-K, Chen F-C, Kuo K-L, Chie W-C, Yang R-S. Sarcopenia screened with SARC-F questionnaire is associated with quality of life and 4-year mortality. *J Am Med Dir Assoc.* 2016;17(12):1129–35. <https://doi.org/10.1016/j.jamda.2016.07.029>.

61. Ida S, Kaneko R, Murata K. SARC-F for screening of sarcopenia among older adults: a Meta-analysis of screening test accuracy. *J Am Med Dir Assoc*. 2018;19(8):685–9. <https://doi.org/10.1016/j.jamda.2018.04.001>.
62. Wang P-y, Li Y, Wang Q. Sarcopenia: an underlying treatment target during the COVID-19 pandemic. *Nutrition*. 2021;84:1111104. <https://doi.org/10.1016/j.nut.2020.1111104>.
63. Gorzelitz J, Farber C, Gangnon R, Cadmus-Bertram L. Accuracy of wearable trackers for measuring moderate-to vigorous-intensity physical activity: a systematic review and Meta-analysis. *Journal for the measurement of physical. Behaviour*. 2020;1(aop):1–12.
64. Kwan RY, Lee D, Lee PH, Tse M, Cheung DS, Thiamwong L, et al. Effects of an mHealth brisk walking intervention on increasing physical activity in older people with cognitive frailty: pilot randomized controlled trial. 2020;8(7):e16596. <https://doi.org/10.2196/16596>.
65. Lai T-F, Liao Y, Lin C-Y, Huang W-C, Hsueh M-C, Chan D-C. Moderate-to-vigorous physical activity duration is more important than timing for physical function in older adults. *Sci Rep*. 2020;10(1):21344. <https://doi.org/10.1038/s41598-020-78072-0>.
66. Beaudart C, Dawson A, Shaw SC, Harvey NC, Kanis JA, Binkley N, et al. Nutrition and physical activity in the prevention and treatment of sarcopenia: systematic review. *Osteoporos Int*. 2017;28(6):1817–33. <https://doi.org/10.1007/s00198-017-3980-9>.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

