

RESEARCH

Open Access



# Changes in objectively measured lifestyle factors during the COVID-19 pandemic in community-dwelling older adults

Takuya Ataka<sup>\*</sup>, Noriyuki Kimura, Atsuko Eguchi and Etsuro Matsubara

## Abstract

**Background:** In this manuscript, we investigate whether objectively measured lifestyle factors, including walking steps, sedentary time, amount of unforced physical activity, level of slight and energetic physical activity, conversation time, and sleep parameters, were altered before and during the COVID-19 pandemic among community-dwelling older adults.

**Methods:** Data were obtained from a prospective cohort study conducted from 2015 to 2019 and a subsequent dementia prevention study undertaken in September 2020. Community-dwelling adults aged  $\geq 65$  years wore wearable sensors before and during the pandemic.

**Results:** A total of 56 adults were enrolled in this study. The mean age was  $74.2 \pm 3.9$  years, and 58.9% ( $n = 33$ ) of the participants were female. Moderate and vigorous physical activity time significantly decreased, and sedentary time significantly increased during the pandemic.

**Conclusions:** This is the first study to demonstrate differences in objectively assessed lifestyle factors before and during the COVID-19 pandemic among community-dwelling older adults. The findings show that the pandemic has adversely affected physical activity among older adults living on their own in Japan.

**Keywords:** Aged, COVID-19, Japan, Lifestyle, Wearable sensor

## Background

The coronavirus (COVID-19) pandemic is threatening the health of human populations globally. Following the emergence of COVID-19 in late 2019, countries experienced the first to sixth waves of the disease. As of March 3, 2022, there have been over 430 million cases of the disease and over 5 million deaths worldwide [1]. The COVID-19 pandemic has forced governments to impose strict confinement rules on their citizens. In Japan, more than 5 million cases of COVID-19 have been confirmed, including approximately 24,000 deaths [2]. The

government declared a state of emergency as the first wave hit, which was not legally binding but prompted many people to wear face masks, practice social distancing, and refrain from going out. The declaration, released on April 16, 2020, applied to the entire country and was in place for more than a month. People restricted their daily activities to prevent the spread of COVID-19 while the rules were in place. These control measures potentially resulted in physical inactivity and social isolation leading to increased levels of psychological stress [3]. Several studies have examined changes in lifestyle factors among young adults that occurred during the period of confinement. The results showed that COVID-19 restrictions had adverse effects on physical activity and sleep in younger adults [4, 5]. However, few studies have

\*Correspondence: t-ataka@oita-u.ac.jp  
Department of Neurology, Faculty of Medicine, Oita University, Oita, Japan



demonstrated whether the COVID-19 pandemic affected the lifestyles of older adults living alone in the community. Physical inactivity, sleep disturbances, and social isolation are major problems in older adults with increasing life prospects [6–9] and are associated with reduced well-being and an increased risk of mental impairment, major preexisting conditions, and death in older adults [10–15]. Thus, comparing lifestyle factors before and during the pandemic is important in understanding their indirect adverse impacts on physical health among older adults. Previously, to examine the risks and protections associated with lifestyle factors in relation to dementia, we conducted a prospective cohort study that objectively measured lifestyle factors in community-dwelling older adults using wearable sensors from August 2015 to March 2019 [16]. Moreover, a subsequent dementia prevention study provided us with an opportunity to measure lifestyle factors objectively in a subgroup of these participants during the pandemic. These noninvasive and cost-effective wearable sensors were used to objectively assess total daily effort and sleep without recall bias, particularly in older adults. Herein, we aimed to evaluate the changes that occurred in objectively measured lifestyle factors before and during the COVID-19 pandemic in community-living older adults.

## Methods

### Study design and participants

A total of 56 participants (33 females, mean age  $74.2 \pm 3.9$  years, years of education  $11.7 \pm 2.0$  years) were enrolled among a population of community-dwelling older adults living in Usuki, a rural city in Oita Prefecture, Japan. The population included both those who live alone and those who do not. In the present study, the inclusion criteria met by all participants were as follows: 1) aged 65 and over, 2) lived in Usuki, 3) physically and psychologically healthy, 4) no dementia, and 5) conducted daily living activities independently. All subjects were instructed to wear wristbands (Silme<sup>™</sup> W20, TDK Corporation Tokyo, Japan) throughout the day, except when showering, for a week during September 2020. These subjects had previously participated in a prospective cohort study with a continuous follow-up that analyzed the association between objectively quantified lifestyle factors and mental function from August 2015 to March 2019. In this study, we defined lifestyle factors as physical movement, sleep habits, and social interaction. During that cohort study, we assessed lifestyle factors using wearable sensors every 3 months for 3 years and calculated average annual data [16]. Overall, the average annual data for 2016 and 2018 and the average weekly data for 2020 were collected for analysis.

### Data obtained by the wearable sensors

For movement detection, we used a tri-axis accelerometer that measures acceleration in three perpendicular axes. Accelerometers are based on the principle of differential capacitance, which results from the movement of the sensing element due to acceleration. Data were recorded continuously and analyzed every minute. We defined nonwearing time as the time when heart rate counts were zero. We analyzed the data for participants who had at least three days of valid accelerometer monitoring during the measurement period and at least three hours of valid accelerometer monitoring per day. We used our wearable sensors to assess various lifestyle parameters, including moving steps, metabolic equivalents of tasks (METs), total sleep time (TST), sleep effectiveness, wake time after sleep onset (WASO), number of awakenings, and conversation time. We summarized the measured values of these parameters into the sum of the sensor data collected on each day and their average values over the entire length of the period. Each parameter was represented as an average daily value. Before our study, we validated the accuracy of our measurements of hiking steps, talking time, and sleep time by comparing them with data from video observations of healthy older subjects.

### Physical activity

We defined walking steps as a movement in the frequency range of 2–3 Hz of acceleration detected by the sensor. The device computed the intensity of activities as METs using algorithms developed by the product designer. Sedentary behavior was defined as activities that involved  $\leq 1.5$  METs such as sitting, lying down, or watching television. Light physical activity (LPA) was defined as activities that involved 1.5–3.0 METs such as slow walking, laundry, cooking food, washing dishes, or vacuuming, whereas moderate and vigorous physical activities (MVPA) were defined as activities that involved  $\geq 3$  METs such as walking, jogging, or ascending and descending stairs.

### Sleep

We evaluated sleep–wake parameters using the magnitude of acceleration and cumulative energy synthesized by a triaxial accelerometer. The data were verified and corrected visually by a qualified technician. Bedtime was determined according to the number of activities logged by the wristband sensor. Sleep factors, including TST, WASO, and sleep efficiency, and the awakening count were measured between 6:00 pm and 5:59 am (the following day). The time of sleep onset was defined as the time when the resting state began, with no movement for

more than 20 min. We evaluated sleep fragmentation by using WASO, sleep effectiveness, and awakening counts. We defined nightly awakenings as 5 to 90 min of continuous movement during a continuous sleep period. Sleep effectiveness was determined as the percentage of TST over bedtime. In this study, sleep diaries were not utilized, rather, the total bedtime was measured using TST and WASO.

### Conversation time

This sensor was able to detect whether an adult or someone nearby had made utterances. The sensor does not record the sound itself but rather the sound as sound pressure and duration. Although we were not able to exclude the speech of someone nearby from the audio data or self-chat, the contribution of the participant in the conversation itself was judged to be a valuable sign of social activity. The microphone on the wearable sensor cannot detect the substance of the chat, but it can collect data as sound every minute. We analyzed sound data to evaluate the duration of the conversation. Our wearable sensor can detect the sound pressure level of utterances within a 2 m radius of the device. The vibration level, considered utterances at this distance, was 55–75 dBA. Furthermore, the incidence band corresponding to the human voice was extracted as a signal frame from the sound data within the vibration range. A chat was defined as a period of one minute during which there were more than four sound frames.

### Statistical analysis

The annual average data for 2016 and 2018 and the weekly average data for 2020 before and during the pandemic, respectively, were used for statistical analyses. A repeated-measures analysis of variance (rANOVA) was conducted to compare nine variables of lifestyle factors (walking steps, sedentary time, LPA, MVPA, chat time, TST, WASO, sleep effectiveness, and waking time count) before and during the pandemic. All statistical analyses were conducted using SPSS statistical software (version 25.0, IBM Corporation, USA) and Prism (version 7.00, GraphPad), and all *p* values of < 0.05 were considered statistically significant.

### Results

The demographic and clinical characteristics of all participants are summarized in Table 1. The mean age was  $74.2 \pm (3.9)$  years, and 41.1% were male ( $n=23$ ). The mean education level was  $11.7 \pm 2.0$  years, and the mean Mini-Mental State Examination score was  $27.4 \pm 2.6$  points. The temporal changes in the objectively measured lifestyle factors are shown in Table 2 and Fig. 1. The amount of daily sedentary time increased ( $p=0.028$ ), and

**Table 1** Clinical and demographical characteristics of participants

Characteristics	
Age, mean (SD), years	74.2 (3.9)
Sex	
Male, n (%)	23 (41.1%)
Female, n (%)	33 (58.9%)
Education level, mean (SD), years	11.7 (2.0)
MMSE, mean (SD)	27.4 (2.6)
Medical history	
Hypertension, n (%)	30 (53.6%)
Diabetes, n (%)	13 (23.2%)
Hypercholesterolemia, n (%)	20 (35.7%)

MMSE Mini-Mental State Examination

the daily MVPA time significantly decreased ( $p=0.042$ ). No significant changes were noticed in daily walking steps, chat time, LPA time, or other sleep parameters.

### Discussion


We investigated the changes in lifestyle factors before and during the COVID-19 pandemic among community-dwelling older adults using wearable sensors. Although multiple reports have assessed the changes in physical activity or sleep parameters individually during the pandemic among younger adults, to the best of our knowledge, this is the first study to investigate the changes in objectively measured lifestyle factors during the pandemic in older adults. Our results provide novel and interesting insights that will help in developing strategies to prevent the indirect health impacts of the pandemic on older adults. Sedentary time significantly increased, while daily MVPA time decreased. The present study has several strengths, including the objective measurement of various lifestyle factors among community-dwelling older adults and the comparison of those measurements obtained before and during the pandemic.

The truly interesting finding of the present study was that the extent of sedentary time was significantly greater during the pandemic than before the pandemic. Few studies have compared the intensity of exercise performed by people before and during the pandemic. One study showed that the time spent undertaking MVPA decreased during a semilockdown period compared with that measured before the pandemic in Chinese young adults [17]. Similarly, our results showed that the daily time spent undertaking MVPA decreased significantly during the pandemic compared with that measured before the pandemic; however, the time spent on MVPA had already decreased between 2016 and 2018. Thus, we were unable to determine whether the decreased

**Table 2** Temporal changes in objectively measured lifestyle factors

	2016	2018	2020	Repeated measures ANOVA p value
Walking steps, mean (SD), steps/day	5727.3 (2506.0)	5449.6 (2547.4)	4985.3 (3092.8)	0.066
Talking ratio, mean (SD), min/day	217.2 (77.8)	222.0 (70.0)	217.1 (78.5)	0.19
TST, mean (SD), min/day	403.1 (66.7)	406.9 (86.2)	400.8 (89.1)	0.67
WASO, mean (SD), min/day	16.4 (9.3)	16.9 (9.6)	17.0 (13.1)	0.44
Sleep efficiency, mean (SD), %/day	96.1 (2.2)	95.9 (2.5)	95.7 (3.4)	0.49
Awake time count, mean (SD), counts/day	0.44 (0.28)	0.43 (0.28)	0.45 (0.33)	0.51
Sedentary time, mean (SD), min/day	785.3 (69.7)	787.2 (72.2)	829.1 (87.4)	0.028
LPA, mean (SD), min/day	26.9 (18.3)	25.5 (15.1)	23.9 (14.4)	0.80
MVPA, mean (SD), min/day	29.4 (16)	26.5 (16)	24.8 (18.5)	0.043

LPA light physical activity, MVPA moderate-vigorous physical activity, TST total sleep time, WASO time awake after sleep



amount of MVPA was caused by aging or the pandemic. Physical inactivity results in increased risks of developing metabolic diseases [18], cardiovascular diseases [19], and immune dysfunctions [20]. For older adults, even short-term physical inactivity can result in muscle loss or frailty [3]. Muscle loss has a negative relationship with metabolic syndromes and cognitive function. The World Health Organization (WHO) has recommended regular physical activity during the pandemic for people of all ages to maintain their health and well-being [21]. Our results suggest that the pandemic indirectly had an adverse impact on physical activity among older adults and support the recommendations of the WHO. Moreover, our results indicated that there were no changes in sleep parameters according to the measurements of older adults taken before and during the pandemic. Previous studies that objectively measured sleep time using smartphones showed that sleep time significantly increased during the pandemic compared with that before the pandemic among young adults [4, 22,

23]. Possible explanations for this discrepancy may have been the differences in the ages of the participants or the severity of psychological stress during the pandemic. The psychological stress endured by Japanese people was milder than that endured by people from other countries because the lockdown in Japan was not as strict.

This study showed no change in conversation time during the pandemic. Few studies have examined the association between conversation time and the pandemic. In this study, we used conversation time as a surrogate marker of social interaction. Conversation potentially includes talking to others, including family members, talking to oneself, and the sound of TV or radio in our method. The decrease in the number of walking steps suggested a possible decrease in interaction with others; however, the conversation time did not actually decrease. It is possible that the time talking to others decreased, while the time spent self-chatting or watching the radio or TV at home increased, but we cannot reveal the detail in this study.

To improve the adverse impacts on physical activity in older people, we recommend the activities suggested by the WHO, such as engaging in exercise online, dancing to music, strength and balance training, and climbing stairs [21]. In addition, we recommend breaking up and standing up regularly while sitting. There were several limitations to the present study that should be considered. First, noise from the television or radio might be detected during the conversation using the chosen detection method, which was based on vibrations and frequencies. In addition, sleep may have been incorrectly categorized as chat time during television broadcasting or listening to the radio. However, a previous study showed that only 6.4% of sleeping time was included in the daily chat. Second, a small number of participants were included in the present study. During a pandemic, it is very difficult to study large populations, but lifestyle factors need to be evaluated in larger populations while maintaining infection control in the future. In summary, this study is the first to confirm the indirect and adverse effects of the COVID-19 pandemic on physical health among community-dwelling older adults. A decline in the amount of daily MVPA time and an increase in sedentary time were observed during the pandemic compared with before the pandemic. These results will contribute to our knowledge about the reduction of physical inactivity during the pandemic in older adults.

#### Acknowledgements

We gratefully acknowledge the assistance of Usuki city employees for their efforts in recruiting adults. We thank Suzuki Co., Ltd., for their assistance with the data collection and HCL Technologies Confidential and Fusa Matsuzaki for the database construction and data analysis. The authors are sincerely grateful to all the participants who enrolled in this study.

#### Disclosure statement

This research was supported by the Japan Agency for Medical Research and Development [Grant Number 18he1402003] and Oita Prefectural Office.

#### Authors' contributions

N.K. had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Concept and design: N.K., E.M.; acquisition, analysis, or interpretation of data: all authors; drafting of the manuscript: T.A., N.K.; critical revision of the manuscript for important intellectual content: all authors; statistical analysis: T.A.; obtained funding: E.M.; and supervision: E.M. All authors have read and approved the manuscript.

#### Funding

The funders had no role in the design or conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or the decision to submit the manuscript for publication.

#### Availability of data and materials

The datasets used during the current study are available from the corresponding author upon reasonable request.

## Declarations

#### Ethics approval and consent to participate

The study was conducted in accordance with the Helsinki Declaration and approved by the institutional review board of Oita University Hospital. All participants provided written informed consent before the study.

#### Consent for publication

Not applicable.

#### Competing interest

Dr. Kimura received a grant from the Japan Society for the Promotion of Science and received honoraria from Takeda Pharmaceutical, Janssen Pharmaceutical, Daiichi Sankyo, Eisai, Sumitomo Dainippon Pharma, FUJIFILM Toyama, Kyowa Kirin, Otsuka Pharmaceutical, and Tsumura outside the submitted work. No other disclosures were reported.

Received: 26 August 2021 Accepted: 1 April 2022

Published online: 14 April 2022

## References

- WHO Coronavirus (COVID-19) Dashboard. World Health Organization. 2021. <https://covid19.who.int/>, Accessed 3 Mar 2022.
- Outbreaks in Japan. Japanese Ministry of Health, Labour and Welfare. 2021. <https://www.mhlw.go.jp/stf/covid-19/kokunainohasseijoukyou.html>, Accessed 3 Mar 2022
- Cavicchioli M, Ferrucci R, Guidetti M, Canevini MP, Pravettoni G, Galli F. What will be the impact of the COVID-19 quarantine on psychological distress? Considerations based on a systematic review of pandemic outbreaks. *Healthcare*. 2021;9:101.
- Kirwan R, McCullough D, Butler T, Perez de Heredia F, Davies IG, Stewart C. Sarcopenia during COVID-19 lockdown restrictions: long-term health effects of short-term muscle loss. *GeroScience*. 2020;42:1547–78.
- Ong JL, Lau T, Massar SA, Chong ZT, Ng BK, Koek D, Zhao W, Yeo BT, Cheong K, Chee MW. COVID-19-related mobility reduction: Heterogeneous effects on sleep and physical activity rhythms. *Sleep*. 2021;44:179.
- Fried LP, Guralnik JM. Disability in older adults: Evidence regarding significance, etiology, and risk. *J Am Geriatr Soc*. 1997;45:92–100.
- Foley DJ, Monjan AA, Brown SL, Simonsick EM, Wallace RB, Blazer DG. Sleep complaints among elderly persons: An epidemiologic study of three communities. *Sleep*. 1995;18:425–32.
- Chien MY, Chen HC. Poor sleep quality is independently associated with physical disability in older adults. *J Clin Sleep Med*. 2015;11:225–32.
- Courtin E, Knapp M. Social isolation, loneliness and health in old age: A scoping review. *Health Soc Care Community*. 2017;25:799–812.
- Buchman AS, Boyle PA, Yu L, Shah RC, Wilson RS, Bennett DA. Total daily physical activity and the risk of AD and cognitive decline in older adults. *Neurology*. 2012;78:1323–9.
- Taylor AH, Cable NT, Faulkner G, Hillsdon M, Narici M, Van Der Bij AK. Physical activity and older adults: A review of health benefits and the effectiveness of interventions. *J Sports Sci*. 2004;22:703–25.
- Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Sleep duration and all-cause mortality: A systematic review and meta-analysis of prospective studies. *Sleep*. 2010;33:585–92.
- Miyata S, Noda A, Iwamoto K, Kawano N, Okuda M, Ozaki N. Poor sleep quality impairs cognitive performance in older adults. *J Sleep Res*. 2013;22:535–41.
- Malcolm M, Frost H, Cowie J. Loneliness and social isolation causal association with health-related lifestyle risk in older adults: a systematic review and meta-analysis protocol. *Syst Rev*. 2019;8:48.
- Steptoe A, Shankar A, Demakakos P, Wardle J. Social isolation, loneliness, and all-cause mortality in older men and women. *Proc Natl Acad Sci U S A*. 2013;110:5797–801.
- Kimura N, Aso Y, Yabuuchi K, Ishibashi M, Hori D, Sasaki Y, Nakamichi A, Uesugi S, Jikumaru M, Sumi K, Eguchi A. Association of modifiable lifestyle factors with cortical amyloid burden and cerebral glucose metabolism in older adults with mild cognitive impairment. *JAMA Netw Open*. 2020;3:e205719.



17. He M, Xian Y, Lv X, He J, Ren Y. Changes in body weight, physical activity, and lifestyle during the semi-lockdown period after the outbreak of COVID-19 in China: An online survey. *Disaster Med Public Health Prep.* 2021;15:e23-28.
18. Martínez-Ferran M, de la Guía-Galipienso F, Sanchis-Gomar F, Pareja-Galeano H. Metabolic impacts of confinement during the COVID-19 pandemic due to modified diet and physical activity habits. *Nutrients.* 2020;12:1549.
19. Wahid A, Manek N, Nichols M, Kelly P, Foster C, Webster P, Kaur A, Friedemann Smith C, Wilkins E, Rayner M, Roberts N, Scarborough P. Quantifying the association between physical activity and cardiovascular disease and diabetes: A systematic review and meta-analysis. *J Am Heart Assoc.* 2016;5:e002495.
20. Weyh C, Krüger K, Strasser B. Physical activity and diet shape the immune system during aging. *Nutrients.* 2020;12:622.
21. #HealthyAtHome - Physical activity. World Health Organization. 2021. <https://www.who.int/news-room/campaigns/connecting-the-world-to-combat-coronavirus/healthyathome/healthyathome---physical-activity#>. Accessed 24 Mar 2022.
22. Sun S, Folarin AA, Ranjan Y, Rashid Z, Conde P, Stewart C, Cummins N, Matcham F, Dalla Costa G, Simblett S, Leocani L. Using smartphones and wearable devices to monitor behavioral changes during COVID-19. *J Med Internet Res.* 2020;22:e19992.
23. Jahrami H, BaHammam AS, Bragazzi NL, Saif Z, Faris M, Vitiello MV. Sleep problems during the COVID-19 pandemic by population: a systematic review and meta-analysis. *J Clin Sleep Med.* 2021;17:299–313.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

