

MEETING ABSTRACT

Open Access

Larynx organ preservation in subjects with hypopharynx-larynx cancer

S Conticello^{1*}, A Fulcheri¹, M G Gorzegno², G Malinverni³, M G Ruo Redda⁴, P Gabriele³, S Aversa¹, C Ondolo¹

From de Senectute: Age and Health Forum
Catanzaro, Italy. 5-7 December 2009

Background

Till not many years ago, the recommended treatment for advanced laryngeal cancer was a total laryngectomy followed by adjuvant radiation therapy. Studies conducted from 1980's have broadened the therapeutic options, in the attempt to cure the cancer while sparing laryngeal functions. Functional surgery and a combination of chemotherapy (CT) and radiotherapy (RT) appears to be able to reach these objectives. We analyzed the case series of patients with laryngeal-hypopharyngeal cancer observed in the period 1999-2006 to evaluate the results on the basis of the treatment offered, focusing in particular on advanced-stage cancer.

Materials and methods

146 patients were observed, of which 137 were analyzed; 113 patients were operated, while 24 had non-surgical treatment: 12 of these had a CT-RT protocol with induction CT followed by concomitant CT-RT.

Results

After surgery in advanced-stage laryngeal cancer survival was 82% at 18 months, and 36% at 5 years, while after CT-RT protocol was 60% at 18 months. There was a statistically relevant advantage in the surgery group ($P=0.02$). For what concerns advanced-stage hypopharyngeal cancer, survival observed was 60% at 3 years (48% at 5 years) in the surgery group, and 62% at 3 years in the CT-RT group, without difference at the log-rank test ($P=0.79$).

Conclusions

In our case history, surgery had a better outcome than CT-RT in advanced-stage larynx cancer, even if we analyzed small groups of patients. These data need to be

confirmed by larger case series; the advancements in surgery, chemotherapy and radiotherapy have furthermore to be held in due consideration, since they are opening interesting perspectives in the treatment of advanced-stage larynx cancer.

Author details

¹Department of Otorhinolaryngology, University of Turin, Azienda Ospedaliera San Luigi Gonzaga, Orbassano (TO), Italy. ²Department of Medical Oncology, University of Turin, Azienda Ospedaliera San Luigi Gonzaga, Orbassano (TO), Italy. ³Department of Radiotherapy, Institute for the Research and the Treatment of Cancer, Candiolo (TO), Italy. ⁴Department of Radiotherapy, University of Turin, Azienda Ospedaliera San Luigi Gonzaga, Orbassano (TO), Italy.

Published: 19 May 2010

doi:10.1186/1471-2318-10-S1-A53

Cite this article as: Conticello et al.: Larynx organ preservation in subjects with hypopharynx-larynx cancer. *BMC Geriatrics* 2010 10(Suppl 1):A53.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

